



L8 ANSWER 7 OF 18 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 1997:164886 CAPLUS

DOCUMENT NUMBER: 126:145606

ORIGINAL REFERENCE NO.: 126:28129a, 28132a

TITLE: Synthesis, Surface Properties, and Hydrolysis of  
Chemodegradable Anionic Surfactants:

## Diastereomerically Pure Sodium *cis*- and *trans*-2-*n*-Alkyl-1,3-dioxan-5-yl Sulfates

AUTHOR(S): Piasecki, Andrzej; Sokołowski, Adam; Burczyk, Bogdan;  
Gancarz, Roman; Kotlewska, Urszula

CORPORATE SOURCE: Institute of Organic and Polymer Technology

Institute of Organic Chemistry Biochemistry and Biotechnology, Technical University of Wrocław, Wrocław, 50-370, Pol.

SOURCE: Langmuir (1997), 13(6)

CODEN: LANGD5; ISSN: 0743-7463

PUBLISHER: *AMERICAN CHEMICAL SOCIETY*

PUBLISHER: American Chemical Society  
DOCUMENT TYPE: Journal

DOCUMENT TYPE:  
LANGUAGE:

LANGUAGE: English  
AB A systematic study concerning

## AB A systematic study concerning and hydrolytic decomposition

and hydrolytic decomposition of pure sodium *cis*- and *trans*-

pure sodium cis- and trans-2-n-alkyl-1,3-dioxan-5-yl sulfates (alkyl: n-C<sub>7</sub>H<sub>15</sub>, n-C<sub>9</sub>H<sub>19</sub>, and n-C<sub>11</sub>H<sub>23</sub>) has been undertaken. Surface parameters of the compds. under study at the aqueous solution/air interface, i.e., surface tension reduction, surface excess concentration, surface area demand per mol.,

and

standard free energy of adsorption and micellization, show differences both in the alkyl chain length and in the hydrophilic, i.e., sulfate, group configuration at the 1,3-dioxane ring. The cmc values are lower for the trans-isomers than for the cis-isomers, the  $\Delta G^\circ_{\text{ads}}$  and  $\Delta G^\circ_{\text{cmc}}$  values are lower for trans-isomers, and the effectiveness of surface tension reduction is higher for the cis-isomers than for the trans-isomers. The investigated compds. undergo an easy hydrolysis reaction of the acetal function, leading to starting aldehydes and sulfated glycerol. The trans-isomers are hydrolyzed much faster than cis-isomers, and no isomerization reaction of the type cis .dblharw. trans is observed during the hydrolysis process.

IT 18445-26-6 18445-27-7

RL: PRP (Properties); RCT (Reactant); RACT (Reactant or reagent)

(intermediate; synthesis, surface properties, and hydrolysis of chemodegradable sodium *cis*- and *trans*-2-*n*-alkyl-1,3-dioxan-5-yl sulfate anionic surfactants)

RN 18445-26-6 CAPLUS

CN 1,3-Dioxan-5-ol, 2-nonyl-, cis- (CA INDEX NAME)

## Relative stereochemistry.

